

## 1 Extended Euclid: Two Ways

In this problem, we will explore the Extended Euclid's Algorithm: first, the traditional implementation, and second, a faster, iterative version. Both ways yield the same result.

Parts (b) and (c) explore the traditional Extended Euclid's Algorithm. The bolded numbers below keep track of which numbers appeared as inputs to the gcd call. Remember that we are interested in writing the GCD as a linear combination of the original inputs, so we don't want to accidentally simplify the expressions and eliminate the inputs.

- (a) As motivation, suppose we've found values of  $a$  and  $b$  such that  $54a + 17b = 1$ . With this knowledge, what is  $17^{-1} \pmod{54}$ ?
- (b) Note that  $x \bmod y$ , by definition, is always  $x$  minus a multiple of  $y$ . So, in the execution of Euclid's algorithm, each newly introduced value can always be expressed as a "combination" of the previous two, like so:

$$\begin{array}{ll}
 \gcd(54, 17) = \gcd(17, 3) & \mathbf{3} = 1 \times \mathbf{54} - 3 \times \mathbf{17} \\
 = \gcd(3, 2) & \mathbf{2} = 1 \times \mathbf{17} - \_\_\_ \times \mathbf{3} \\
 = \gcd(2, 1) & \mathbf{1} = 1 \times \mathbf{3} - \_\_\_ \times \mathbf{2} \\
 = \gcd(1, 0) & [\mathbf{0} = 1 \times \mathbf{2} - \_\_\_ \times \mathbf{1}] \\
 = 1. &
 \end{array}$$

(Fill in the blanks)

- (c) Recall that our goal is to fill out the blanks in

$$1 = \_\_\_ \times \mathbf{54} + \_\_\_ \times \mathbf{17}.$$

To do so, we work back up from the bottom, and express the gcd above as a combination of the two arguments on each of the previous lines:

$$\begin{aligned}
 1 &= \_\_\_ \times \mathbf{3} + \_\_\_ \times \mathbf{2} \\
 &= \\
 &= \_\_\_ \times \mathbf{17} + \_\_\_ \times \mathbf{3} \\
 &= \\
 &= \_\_\_ \times \mathbf{54} + \_\_\_ \times \mathbf{17}
 \end{aligned}$$

What does this imply, in this case, about the multiplicative inverse of 17, in arithmetic mod 54?

- (d) In the previous parts, we used a recursive method to write  $\gcd(54, 17)$  as a linear combination of 54 and 17. We can also compute the same result iteratively—this is an alternative to the above method that is oftentimes faster. We begin by writing equations for our initial arguments, 54 and 17, as a linear combination of themselves:

$$54 = 1 \times 54 + 0 \times 17 \quad (E_1)$$

$$17 = 0 \times 54 + 1 \times 17 \quad (E_2)$$

We can then use these initial equations (labeled  $E_1$  and  $E_2$  for ease of reference) to iteratively write reduced values as linear combinations of 54 and 17, until we are able to write an equation for  $\gcd(54, 17)$ , as desired.

In particular, we want to subtract as many multiples of the second equation as possible from the first to create a new equation with a lower LHS value. We can keep iterating until the LHS becomes  $\gcd(54, 17) = 1$ .

$$\text{---} = \text{---} \times 54 + \text{---} \times 17 \quad (E_3 = E_1 - \text{---} \times E_2)$$

$$\text{---} = \text{---} \times 54 + \text{---} \times 17 \quad (E_4 = E_2 - \text{---} \times E_3)$$

$$1 = \text{---} \times 54 + \text{---} \times 17 \quad (E_5 = E_3 - \text{---} \times E_4)$$

What does this imply, in this case, about the multiplicative inverse of 17, in arithmetic mod 54? Verify that your answer is equivalent to the previous part.

- (e) Calculate the gcd of 17 and 39, and determine how to express this as a “combination” of 17 and 39. What does this imply, in this case, about the multiplicative inverse of 17, in arithmetic mod 39?

## 2 Unique Prime Factorization

We proved in lecture that every positive integer has a factorization into primes. Using the techniques in this course, we can show that this factorization is unique, up to reordering the factors!

Recall from lecture that we defined a prime number  $p$  to be a positive integer whose only positive factors are 1 and  $p$ .

**In this problem, you should not assume that the positive integers have unique prime factorization.**

- (a) Let  $p$  and  $q$  be distinct primes. Show that  $p \nmid q$ .

(b) Prove that if  $a$  and  $b$  are positive integers such that  $p \mid ab$ , then  $p \mid a$  or  $p \mid b$ . (Hint: Use the Extended Euclidean algorithm: There are integers  $x, y$  such that  $ax + py = \gcd(a, p)$ .)

(c) Show that the prime factorization of a positive integer  $n$  is unique up to reordering the factors. (Hint: Suppose that there were two prime factorizations of  $n$ . Use the previous parts to show that the factorizations are the same.)

### 3 Chinese Remainder Theorem Practice

In this question, you will solve for a natural number  $x$  such that,

$$\begin{aligned}x &\equiv 1 \pmod{3} \\x &\equiv 3 \pmod{7} \\x &\equiv 4 \pmod{11}\end{aligned}\tag{1}$$

(a) Suppose you find 3 natural numbers  $a, b, c$  that satisfy the following properties:

$$a \equiv 1 \pmod{3}; a \equiv 0 \pmod{7}; a \equiv 0 \pmod{11},\tag{2}$$

$$b \equiv 0 \pmod{3}; b \equiv 3 \pmod{7}; b \equiv 0 \pmod{11},\tag{3}$$

$$c \equiv 0 \pmod{3}; c \equiv 0 \pmod{7}; c \equiv 4 \pmod{11}.\tag{4}$$

Show how you can use the knowledge of  $a, b$  and  $c$  to compute an  $x$  that satisfies (1).

In the following parts, you will compute natural numbers  $a, b$  and  $c$  that satisfy the above 3 conditions and use them to find an  $x$  that satisfies (1).

- (b) Find a natural number  $a$  that satisfies (2). That is,  $a \equiv 1 \pmod{3}$  and is a multiple of 7 and 11.

It may help to start with a number that is a multiple of both 7 and 11; what number should we multiply this by in order to make it equivalent to  $1 \pmod{3}$ ? Try to set up an equation to solve for this quantity.

- (c) Find a natural number  $b$  that satisfies (3). That is,  $b \equiv 3 \pmod{7}$  and is a multiple of 3 and 11.

- (d) Find a natural number  $c$  that satisfies (4). That is,  $c \equiv 4 \pmod{11}$  and is a multiple of 3 and 7.

- (e) Putting together your answers for parts (a), (b), (c) and (d), report an  $x$  that satisfies (1).